

Ce₂O₂S Structure: A2B2C_hP5_164_d_d_a-002


This structure previously used the label(s) **A2B2C_hP5_164_d_d_a**. Calls to these addresses will be redirected here.

Cite this page as:

- H. Eckert, S. Divilov, M. J. Mehl, D. Hicks, A. C. Zettl, M. Esters, X. Campilongo, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 4*, Comput. Mater. Sci. **240**, 112988 (2024), doi: 10.1016/j.commatsci.2024.112988.
- D. Hicks, M. J. Mehl, M. Esters, C. Oses, O. Levy, G. L. W. Hart, C. Toher, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 3*, Comput. Mater. Sci. **199**, 110450 (2021), doi: 10.1016/j.commatsci.2021.110450.

<https://aflow.org/prototype-encyclopedia/FG79>

https://aflow.org/prototype-encyclopedia/A2B2C_hP5_164_d_d_a-002



Prototype	Ce ₂ O ₂ S
AFLOW prototype label	A2B2C_hP5_164_d_d_a-002
ICSD	31639
CCDC	1605642
Pearson symbol	hP5
Space group number	164
Space group symbol	$P\bar{3}m1$

AFLOW prototype command `aflow --proto=A2B2C_hP5_164_d_d_a-002`
 `--params=a, c/a, z2, z3`


Other compounds with this structure

BaCd₂As₂, CaAl₂Si₂, CaAs₂Bi₂, CaAs₂Mn₂, CaAs₂P₂, CaAs₂Sb₂, CaBi₂Mg₂, CaBi₂Mn₂, CaCd₂As₂, CaCd₂P₂, CaMg₂Bi₂, CaMg₂Sb₂, CaMn₂P₂, CaZn₂As₂, CaZn₂P₂, CaZn₂Sb₂, CeCuZnP₂, DyCuZnP₂, ErCuZnP₂, EuAs₂Cd₂, EuBi₂Mg₂, EuCd₂P₂, EuCd₂Sb₂, EuP₂iZn₂, GdCuZnP₂, HoCuZnP₂, LaCuZnP₂, LuCuZnP₂, NdCuZnP₂, PrCuZnP₂, SCe₂O₂, SCe₂Se₂, SLa₂O₂, SO₂Pu₂, ScCuZnP₂, SmCuZnP₂, SrAs₂Mn₂, SrAs₂P₂, SrAs₂Sb₂, SrCd₂As₂, SrCd₂P₂, SrMn₂P₂, TbCuZnP₂, TmCuZnP₂, YCuZnP₂, YbCuZnP₂, YbMg₂Bi₂, YbMnCuP₂

- This is the ternary form of the D_{5h} La₂O₃ and the D_{5h} Al₃Ni₂ structures. We have separated it from the parents because of the large number of compounds involved.
 - Authors after (Zachariasen, 1949) often use CaAl₂Si₂ as the prototype for this structure.
 - If we add an atom to the (1b) site we obtain either the ternary LaLi₃Sb₂ or the quaternary TbLiCu₂P₂ structure.
-

Trigonal (Hexagonal) primitive vectors

$$\begin{aligned}\mathbf{a}_1 &= \frac{1}{2}a\hat{\mathbf{x}} - \frac{\sqrt{3}}{2}a\hat{\mathbf{y}} \\ \mathbf{a}_2 &= \frac{1}{2}a\hat{\mathbf{x}} + \frac{\sqrt{3}}{2}a\hat{\mathbf{y}} \\ \mathbf{a}_3 &= c\hat{\mathbf{z}}\end{aligned}$$



Basis vectors

	Lattice coordinates		Cartesian coordinates	Wyckoff position	Atom type
$\mathbf{B}_1$	$=$	0	$=$	0	(1a) S I
$\mathbf{B}_2$	$=$	$\frac{1}{3}\mathbf{a}_1 + \frac{2}{3}\mathbf{a}_2 + z_2\mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + cz_2\hat{\mathbf{z}}$	(2d) Ce I
$\mathbf{B}_3$	$=$	$\frac{2}{3}\mathbf{a}_1 + \frac{1}{3}\mathbf{a}_2 - z_2\mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} - cz_2\hat{\mathbf{z}}$	(2d) Ce I
$\mathbf{B}_4$	$=$	$\frac{1}{3}\mathbf{a}_1 + \frac{2}{3}\mathbf{a}_2 + z_3\mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} + cz_3\hat{\mathbf{z}}$	(2d) O I
$\mathbf{B}_5$	$=$	$\frac{2}{3}\mathbf{a}_1 + \frac{1}{3}\mathbf{a}_2 - z_3\mathbf{a}_3$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a\hat{\mathbf{y}} - cz_3\hat{\mathbf{z}}$	(2d) O I

References

- [1] W. H. Zachariasen, *Crystal chemical studies of the 5f-series of elements. VII. The crystal structure of Ce₂O₂S, La₂O₂S and Pu₂O₂S*, Acta Cryst. **2**, 60–62 (1949), doi:10.1107/S0365110X49000138.

First cited in

D. Hicks, M. J. Mehl, M. Esters, C. Osos, O. Levy, G. L. W. Hart, C. Toher, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 3*, Comput. Mater. Sci. **199**, 110450 (2021), doi: 10.1016/j.commatsci.2021.110450.