

K₂S₂O₅ (*K*0₁) Structure: A2B5C2_mP18_11_2e_e2f_2e-001


This structure previously used the label(s) **A2B5C2_mP18_11_2e_e2f_2e**. Calls to these addresses will be redirected here.

Cite this page as:

- H. Eckert, S. Divilov, M. J. Mehl, D. Hicks, A. C. Zettel, M. Esters, X. Campilongo, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 4*, Comput. Mater. Sci. **240**, 112988 (2024), doi: 10.1016/j.commatsci.2024.112988.
- D. Hicks, M. J. Mehl, M. Esters, C. Oses, O. Levy, G. L. W. Hart, C. Toher, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 3*, Comput. Mater. Sci. **199**, 110450 (2021), doi: 10.1016/j.commatsci.2021.110450.

<https://afLOW.org/prototype-encyclopedia/YYGA>

https://afLOW.org/prototype-encyclopedia/A2B5C2_mP18_11_2e_e2f_2e-001




Prototype	K ₂ O ₅ S ₂
AFLOW prototype label	A2B5C2_mP18_11_2e_e2f_2e-001
Strukturbericht designation	<i>K</i> 0 ₁
ICSD	60025
CCDC	1623781
Pearson symbol	mP18
Space group number	11
Space group symbol	<i>P</i> 2 ₁ / <i>m</i>
AFLOW prototype command	<pre>afLOW --proto=A2B5C2_mP18_11_2e_e2f_2e-001 --params=a,b/a,c/a,β,x₁,z₁,x₂,z₂,x₃,z₃,x₄,z₄,x₅,z₅,x₆,y₆,z₆,x₇,y₇,z₇</pre>

Other compounds with this structure

$K_2S_2P_5$, $(NH_4)_2S_2P_5$

Simple Monoclinic primitive vectors

$$\begin{aligned} \mathbf{a}_1 &= a \hat{\mathbf{x}} \\ \mathbf{a}_2 &= b \hat{\mathbf{y}} \\ \mathbf{a}_3 &= c \cos \beta \hat{\mathbf{x}} + c \sin \beta \hat{\mathbf{z}} \end{aligned}$$



Basis vectors

	Lattice coordinates		Cartesian coordinates	Wyckoff position	Atom type
$\mathbf{B}_1$	$= x_1 \mathbf{a}_1 + \frac{1}{4} \mathbf{a}_2 + z_1 \mathbf{a}_3$	$=$	$(ax_1 + cz_1 \cos \beta) \hat{\mathbf{x}} + \frac{1}{4}b \hat{\mathbf{y}} + cz_1 \sin \beta \hat{\mathbf{z}}$	(2e)	K I
$\mathbf{B}_2$	$= -x_1 \mathbf{a}_1 + \frac{3}{4} \mathbf{a}_2 - z_1 \mathbf{a}_3$	$=$	$-(ax_1 + cz_1 \cos \beta) \hat{\mathbf{x}} + \frac{3}{4}b \hat{\mathbf{y}} - cz_1 \sin \beta \hat{\mathbf{z}}$	(2e)	K I
$\mathbf{B}_3$	$= x_2 \mathbf{a}_1 + \frac{1}{4} \mathbf{a}_2 + z_2 \mathbf{a}_3$	$=$	$(ax_2 + cz_2 \cos \beta) \hat{\mathbf{x}} + \frac{1}{4}b \hat{\mathbf{y}} + cz_2 \sin \beta \hat{\mathbf{z}}$	(2e)	K II
$\mathbf{B}_4$	$= -x_2 \mathbf{a}_1 + \frac{3}{4} \mathbf{a}_2 - z_2 \mathbf{a}_3$	$=$	$-(ax_2 + cz_2 \cos \beta) \hat{\mathbf{x}} + \frac{3}{4}b \hat{\mathbf{y}} - cz_2 \sin \beta \hat{\mathbf{z}}$	(2e)	K II
$\mathbf{B}_5$	$= x_3 \mathbf{a}_1 + \frac{1}{4} \mathbf{a}_2 + z_3 \mathbf{a}_3$	$=$	$(ax_3 + cz_3 \cos \beta) \hat{\mathbf{x}} + \frac{1}{4}b \hat{\mathbf{y}} + cz_3 \sin \beta \hat{\mathbf{z}}$	(2e)	O I
$\mathbf{B}_6$	$= -x_3 \mathbf{a}_1 + \frac{3}{4} \mathbf{a}_2 - z_3 \mathbf{a}_3$	$=$	$-(ax_3 + cz_3 \cos \beta) \hat{\mathbf{x}} + \frac{3}{4}b \hat{\mathbf{y}} - cz_3 \sin \beta \hat{\mathbf{z}}$	(2e)	O I
$\mathbf{B}_7$	$= x_4 \mathbf{a}_1 + \frac{1}{4} \mathbf{a}_2 + z_4 \mathbf{a}_3$	$=$	$(ax_4 + cz_4 \cos \beta) \hat{\mathbf{x}} + \frac{1}{4}b \hat{\mathbf{y}} + cz_4 \sin \beta \hat{\mathbf{z}}$	(2e)	S I
$\mathbf{B}_8$	$= -x_4 \mathbf{a}_1 + \frac{3}{4} \mathbf{a}_2 - z_4 \mathbf{a}_3$	$=$	$-(ax_4 + cz_4 \cos \beta) \hat{\mathbf{x}} + \frac{3}{4}b \hat{\mathbf{y}} - cz_4 \sin \beta \hat{\mathbf{z}}$	(2e)	S I
$\mathbf{B}_9$	$= x_5 \mathbf{a}_1 + \frac{1}{4} \mathbf{a}_2 + z_5 \mathbf{a}_3$	$=$	$(ax_5 + cz_5 \cos \beta) \hat{\mathbf{x}} + \frac{1}{4}b \hat{\mathbf{y}} + cz_5 \sin \beta \hat{\mathbf{z}}$	(2e)	S II
$\mathbf{B}_{10}$	$= -x_5 \mathbf{a}_1 + \frac{3}{4} \mathbf{a}_2 - z_5 \mathbf{a}_3$	$=$	$-(ax_5 + cz_5 \cos \beta) \hat{\mathbf{x}} + \frac{3}{4}b \hat{\mathbf{y}} - cz_5 \sin \beta \hat{\mathbf{z}}$	(2e)	S II
$\mathbf{B}_{11}$	$= x_6 \mathbf{a}_1 + y_6 \mathbf{a}_2 + z_6 \mathbf{a}_3$	$=$	$(ax_6 + cz_6 \cos \beta) \hat{\mathbf{x}} + by_6 \hat{\mathbf{y}} + cz_6 \sin \beta \hat{\mathbf{z}}$	(4f)	O II
$\mathbf{B}_{12}$	$= -x_6 \mathbf{a}_1 + (y_6 + \frac{1}{2}) \mathbf{a}_2 - z_6 \mathbf{a}_3$	$=$	$-(ax_6 + cz_6 \cos \beta) \hat{\mathbf{x}} + b(y_6 + \frac{1}{2}) \hat{\mathbf{y}} - cz_6 \sin \beta \hat{\mathbf{z}}$	(4f)	O II
$\mathbf{B}_{13}$	$= -x_6 \mathbf{a}_1 - y_6 \mathbf{a}_2 - z_6 \mathbf{a}_3$	$=$	$-(ax_6 + cz_6 \cos \beta) \hat{\mathbf{x}} - by_6 \hat{\mathbf{y}} - cz_6 \sin \beta \hat{\mathbf{z}}$	(4f)	O II
$\mathbf{B}_{14}$	$= x_6 \mathbf{a}_1 - (y_6 - \frac{1}{2}) \mathbf{a}_2 + z_6 \mathbf{a}_3$	$=$	$(ax_6 + cz_6 \cos \beta) \hat{\mathbf{x}} - b(y_6 - \frac{1}{2}) \hat{\mathbf{y}} + cz_6 \sin \beta \hat{\mathbf{z}}$	(4f)	O II
$\mathbf{B}_{15}$	$= x_7 \mathbf{a}_1 + y_7 \mathbf{a}_2 + z_7 \mathbf{a}_3$	$=$	$(ax_7 + cz_7 \cos \beta) \hat{\mathbf{x}} + by_7 \hat{\mathbf{y}} + cz_7 \sin \beta \hat{\mathbf{z}}$	(4f)	O III
$\mathbf{B}_{16}$	$= -x_7 \mathbf{a}_1 + (y_7 + \frac{1}{2}) \mathbf{a}_2 - z_7 \mathbf{a}_3$	$=$	$-(ax_7 + cz_7 \cos \beta) \hat{\mathbf{x}} + b(y_7 + \frac{1}{2}) \hat{\mathbf{y}} - cz_7 \sin \beta \hat{\mathbf{z}}$	(4f)	O III
$\mathbf{B}_{17}$	$= -x_7 \mathbf{a}_1 - y_7 \mathbf{a}_2 - z_7 \mathbf{a}_3$	$=$	$-(ax_7 + cz_7 \cos \beta) \hat{\mathbf{x}} - by_7 \hat{\mathbf{y}} - cz_7 \sin \beta \hat{\mathbf{z}}$	(4f)	O III
$\mathbf{B}_{18}$	$= x_7 \mathbf{a}_1 - (y_7 - \frac{1}{2}) \mathbf{a}_2 + z_7 \mathbf{a}_3$	$=$	$(ax_7 + cz_7 \cos \beta) \hat{\mathbf{x}} - b(y_7 - \frac{1}{2}) \hat{\mathbf{y}} + cz_7 \sin \beta \hat{\mathbf{z}}$	(4f)	O III

References

- [1] I.-C. Chen and Y. Wang, *Reinvestigation of potassium pyrosulfite, $K_2S_2O_5$* **40**, 1780–1781 (1984), doi:10.1107/S0108270184009525.

First cited in

D. Hicks, M. J. Mehl, M. Esters, C. Oses, O. Levy, G. L. W. Hart, C. Toher, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 3*, Comput. Mater. Sci. **199**, 110450 (2021), doi: 10.1016/j.commatsci.2021.110450.