

PI₃ Structure: A3B_hP8_173_c-b-001


This structure previously used the label(s) **A3B_hP8_173_c-b**. Calls to these addresses will be redirected here.

Cite this page as:

- H. Eckert, S. Divilov, M. J. Mehl, D. Hicks, A. C. Zettel, M. Esters, X. Campilongo, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 4*, Comput. Mater. Sci. **240**, 112988 (2024), doi: 10.1016/j.commatsci.2024.112988.
- D. Hicks, M. J. Mehl, M. Esters, C. Osos, O. Levy, G. L. W. Hart, C. Toher, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 3*, Comput. Mater. Sci. **199**, 110450 (2021), doi: 10.1016/j.commatsci.2021.110450.

<https://aflow.org/prototype-encyclopedia/RB2M>

https://aflow.org/prototype-encyclopedia/A3B_hP8_173_c-b-001



Prototype

I₃P

AFLOW prototype label

A3B_hP8_173_c-b-001

ICSD

311

CCDC

1590858

Pearson symbol hP8

Space group number 173

Space group symbol $P6_3$

AFLOW prototype command `aflow --proto=A3B_hP8_173_c_b-001
--params=a, c/a, z1, x2, y2, z2`


Other compounds with this structure

HCl₃

- Space group $P6_3$ #173 allows an arbitrary choice for the origin of the z -axis. We use this to set $z_1 = 0$ for the potassium site.

Hexagonal primitive vectors

$$\begin{aligned}\mathbf{a}_1 &= \frac{1}{2}a \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}a \hat{\mathbf{y}} \\ \mathbf{a}_2 &= \frac{1}{2}a \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}a \hat{\mathbf{y}} \\ \mathbf{a}_3 &= c \hat{\mathbf{z}}\end{aligned}$$



Basis vectors

	Lattice coordinates		Cartesian coordinates		Wyckoff position	Atom type
$\mathbf{B}_1$	$= \frac{1}{3} \mathbf{a}_1 + \frac{2}{3} \mathbf{a}_2 + z_1 \mathbf{a}_3$	$=$	$\frac{1}{2} a \hat{\mathbf{x}} + \frac{\sqrt{3}}{6} a \hat{\mathbf{y}} + cz_1 \hat{\mathbf{z}}$	(2b)		P I
$\mathbf{B}_2$	$= \frac{2}{3} \mathbf{a}_1 + \frac{1}{3} \mathbf{a}_2 + \left(z_1 + \frac{1}{2}\right) \mathbf{a}_3$	$=$	$\frac{1}{2} a \hat{\mathbf{x}} - \frac{\sqrt{3}}{6} a \hat{\mathbf{y}} + c \left(z_1 + \frac{1}{2}\right) \hat{\mathbf{z}}$	(2b)		P I
$\mathbf{B}_3$	$= x_2 \mathbf{a}_1 + y_2 \mathbf{a}_2 + z_2 \mathbf{a}_3$	$=$	$\frac{1}{2} a (x_2 + y_2) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2} a (x_2 - y_2) \hat{\mathbf{y}} + cz_2 \hat{\mathbf{z}}$	(6c)		I I
$\mathbf{B}_4$	$= -y_2 \mathbf{a}_1 + (x_2 - y_2) \mathbf{a}_2 + z_2 \mathbf{a}_3$	$=$	$\frac{1}{2} a (x_2 - 2y_2) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2} ax_2 \hat{\mathbf{y}} + cz_2 \hat{\mathbf{z}}$	(6c)		I I
$\mathbf{B}_5$	$= -(x_2 - y_2) \mathbf{a}_1 - x_2 \mathbf{a}_2 + z_2 \mathbf{a}_3$	$=$	$-\frac{1}{2} a (2x_2 - y_2) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2} ay_2 \hat{\mathbf{y}} + cz_2 \hat{\mathbf{z}}$	(6c)		I I
$\mathbf{B}_6$	$= -x_2 \mathbf{a}_1 - y_2 \mathbf{a}_2 + \left(z_2 + \frac{1}{2}\right) \mathbf{a}_3$	$=$	$-\frac{1}{2} a (x_2 + y_2) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2} a (x_2 - y_2) \hat{\mathbf{y}} + c \left(z_2 + \frac{1}{2}\right) \hat{\mathbf{z}}$	(6c)		I I
$\mathbf{B}_7$	$= y_2 \mathbf{a}_1 - (x_2 - y_2) \mathbf{a}_2 + \left(z_2 + \frac{1}{2}\right) \mathbf{a}_3$	$=$	$\frac{1}{2} a (-x_2 + 2y_2) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2} ax_2 \hat{\mathbf{y}} + c \left(z_2 + \frac{1}{2}\right) \hat{\mathbf{z}}$	(6c)		I I
$\mathbf{B}_8$	$= (x_2 - y_2) \mathbf{a}_1 + x_2 \mathbf{a}_2 + \left(z_2 + \frac{1}{2}\right) \mathbf{a}_3$	$=$	$\frac{1}{2} a (2x_2 - y_2) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2} ay_2 \hat{\mathbf{y}} + c \left(z_2 + \frac{1}{2}\right) \hat{\mathbf{z}}$	(6c)		I I

References

- [1] E. T. Lance, J. M. Haschke, and D. R. Peacor, *Crystal and molecular structure of phosphorus triiodide*, Inorg. Chem. **15**, 780–781 (1976), doi:10.1021/ic50158a007.

Found in

- [1] P. Villars and K. Cenzual, *Pearson's Crystal Data – Crystal Structure Database for Inorganic Compounds* (2013). ASM International.

First cited in

D. Hicks, M. J. Mehl, E. Gossett, C. Toher, O. Levy, R. M. Hanson, G. Hart, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 2*, Comput. Mater. Sci. **161**, S1 (2019). doi: 10.1016/j.commatsci.2018.10.043