


β_1 -K₂UF₆ Structure: A6B2C_hP9_189_fg_c_b-001

Cite this page as:

- H. Eckert, S. Divilov, M. J. Mehl, D. Hicks, A. C. Zettel, M. Esters, X. Campilongo, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 4*, Comput. Mater. Sci. **240**, 112988 (2024), doi: 10.1016/j.commatsci.2024.112988.
- D. Hicks, M. J. Mehl, M. Esters, C. Osos, O. Levy, G. L. W. Hart, C. Toher, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 3*, Comput. Mater. Sci. **199**, 110450 (2021), doi: 10.1016/j.commatsci.2021.110450.

<https://aflow.org/prototype-encyclopedia/XTDJ>

https://aflow.org/prototype-encyclopedia/A6B2C_hP9_189_fg_c_b-001



Prototype	F ₆ K ₂ U
AFLOW prototype label	A6B2C_hP9_189_fg_c_b-001
ICSD	26193
CCDC	1601624
Pearson symbol	hP9
Space group number	189
Space group symbol	$P\bar{6}2m$

AFLOW prototype command `aflow --proto=A6B2C_hP9_189_fg_c_b-001`
 `--params=a, c/a, x3, x4`


Other compounds with this structure

K₂CeF₆, K₂ThF₆, δ -Na₂UF₆

- α -K₂UF₆ is in the cubic fluorite (*C*1, CaF₂) structure, with the potassium and uranium atoms placed randomly on the calcium sites.
- β_2 -K₂UF₆ takes on the trigonal β -Na₂ThF₆ structure.

Hexagonal primitive vectors

$$\begin{aligned}\mathbf{a}_1 &= \frac{1}{2}a\hat{\mathbf{x}} - \frac{\sqrt{3}}{2}a\hat{\mathbf{y}} \\ \mathbf{a}_2 &= \frac{1}{2}a\hat{\mathbf{x}} + \frac{\sqrt{3}}{2}a\hat{\mathbf{y}} \\ \mathbf{a}_3 &= c\hat{\mathbf{z}}\end{aligned}$$



Basis vectors

	Lattice coordinates		Cartesian coordinates	Wyckoff position	Atom type
$\mathbf{B}_1$	$= \frac{1}{2}\mathbf{a}_3$	$=$	$\frac{1}{2}c\hat{\mathbf{z}}$	(1b)	U I
$\mathbf{B}_2$	$= \frac{1}{3}\mathbf{a}_1 + \frac{2}{3}\mathbf{a}_2$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a\hat{\mathbf{y}}$	(2c)	K I
$\mathbf{B}_3$	$= \frac{2}{3}\mathbf{a}_1 + \frac{1}{3}\mathbf{a}_2$	$=$	$\frac{1}{2}a\hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a\hat{\mathbf{y}}$	(2c)	K I
$\mathbf{B}_4$	$= x_3\mathbf{a}_1$	$=$	$\frac{1}{2}ax_3\hat{\mathbf{x}} - \frac{\sqrt{3}}{2}ax_3\hat{\mathbf{y}}$	(3f)	F I
$\mathbf{B}_5$	$= x_3\mathbf{a}_2$	$=$	$\frac{1}{2}ax_3\hat{\mathbf{x}} + \frac{\sqrt{3}}{2}ax_3\hat{\mathbf{y}}$	(3f)	F I
$\mathbf{B}_6$	$= -x_3\mathbf{a}_1 - x_3\mathbf{a}_2$	$=$	$-ax_3\hat{\mathbf{x}}$	(3f)	F I
$\mathbf{B}_7$	$= x_4\mathbf{a}_1 + \frac{1}{2}\mathbf{a}_3$	$=$	$\frac{1}{2}ax_4\hat{\mathbf{x}} - \frac{\sqrt{3}}{2}ax_4\hat{\mathbf{y}} + \frac{1}{2}c\hat{\mathbf{z}}$	(3g)	F II
$\mathbf{B}_8$	$= x_4\mathbf{a}_2 + \frac{1}{2}\mathbf{a}_3$	$=$	$\frac{1}{2}ax_4\hat{\mathbf{x}} + \frac{\sqrt{3}}{2}ax_4\hat{\mathbf{y}} + \frac{1}{2}c\hat{\mathbf{z}}$	(3g)	F II
$\mathbf{B}_9$	$= -x_4\mathbf{a}_1 - x_4\mathbf{a}_2 + \frac{1}{2}\mathbf{a}_3$	$=$	$-ax_4\hat{\mathbf{x}} + \frac{1}{2}c\hat{\mathbf{z}}$	(3g)	F II

References

- [1] G. Brunton, *Refinement of the crystal structure of β_1 -K₂UF₆*, Acta Crystallogr. Sect. B **25**, 2163–2164 (1969), doi:10.1107/S0567740869005310.

Found in

- [1] A. Grzechnik, C. C. Underwood, J. W. Kolis, and K. Frieze, *Crystal structures and stability of K₂ThF₆ and K₇Th₆F₃₁ on compression*, J. Fluor. Chem. **150**, 8–13 (2013), doi:10.1016/j.jfluchem.2013.02.024.

First cited in

H. Eckert, S. Divilov, M. J. Mehl, D. Hicks, A. C. Zettel, M. Esters, X. Campilongo, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 4*, Comput. Mater. Sci. **240**, 112988 (2024), doi: 10.1016/j.commatsci.2024.112988.