

# Cs<sub>3</sub>Tl<sub>2</sub>Cl<sub>9</sub> (*K*7<sub>2</sub>) Structure: A9B3C2\_hR28\_167\_ef\_e\_c-001

This structure previously used the label(s) **A9B3C2\_hR28\_167\_ef\_e\_c**. Calls to these addresses will be redirected here.

Cite this page as:

- H. Eckert, S. Divilov, M. J. Mehl, D. Hicks, A. C. Zettel, M. Esters, X. Campilongo, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 4*, Comput. Mater. Sci. **240**, 112988 (2024), doi: 10.1016/j.commatsci.2024.112988.
- D. Hicks, M. J. Mehl, M. Esters, C. Osos, O. Levy, G. L. W. Hart, C. Toher, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 3*, Comput. Mater. Sci. **199**, 110450 (2021), doi: 10.1016/j.commatsci.2021.110450.

<https://aflow.org/prototype-encyclopedia/7M03>

[https://aflow.org/prototype-encyclopedia/A9B3C2\\_hR28\\_167\\_ef\\_e\\_c-001](https://aflow.org/prototype-encyclopedia/A9B3C2_hR28_167_ef_e_c-001)

|                                    |                                                                                                                                  |
|------------------------------------|----------------------------------------------------------------------------------------------------------------------------------|
| Prototype                          | Cl <sub>9</sub> Cs <sub>3</sub> Tl <sub>2</sub>                                                                                  |
| AFLOW prototype label              | A9B3C2_hR28_167_ef_e_c-001                                                                                                       |
| <i>Strukturbericht</i> designation | <i>K</i> 7 <sub>2</sub>                                                                                                          |
| ICSD                               | 27849                                                                                                                            |
| CCDC                               | 1603027                                                                                                                          |
| Pearson symbol                     | hR28                                                                                                                             |
| Space group number                 | 167                                                                                                                              |
| Space group symbol                 | $R\bar{3}c$                                                                                                                      |
| AFLOW prototype command            | <code>aflow --proto=A9B3C2_hR28_167_ef_e_c-001</code><br><code>--params=<math>a, c/a, x_1, x_2, x_3, x_4, y_4, z_4</math></code> |


## Other compounds with this structure







Cs<sub>3</sub>Dy<sub>2</sub>Br<sub>9</sub>, Cs<sub>3</sub>Er<sub>2</sub>Br<sub>9</sub>, Cs<sub>3</sub>Ho<sub>2</sub>Br<sub>9</sub>, Cs<sub>3</sub>Lu<sub>2</sub>Cl<sub>9</sub>, Cs<sub>3</sub>Tb<sub>2</sub>Br<sub>9</sub>, Cs<sub>3</sub>Yb<sub>2</sub>Br<sub>9</sub>, Ba<sub>3</sub>Os<sub>2</sub>O<sub>9</sub>, Ba<sub>3</sub>W<sub>2</sub>O<sub>9</sub>

- (Hoard, 1935) followed by (Downs, 2003), give the atomic coordinates in the style of (Wyckoff, 1922), who lists space group  $D_{3d}^6$  (the Schönflies notation for space group  $R\bar{3}c$ ) as space group #203, instead of #167, and uses an origin which corresponds to  $(1/4 \ 1/4 \ 1/4)$  in our lattice coordinates. We used FINDSYM to convert this into our standard setting for space group #167.

## Rhombohedral primitive vectors

$$\begin{aligned}\mathbf{a}_1 &= \frac{1}{2}a \hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a \hat{\mathbf{y}} + \frac{1}{3}c \hat{\mathbf{z}} \\ \mathbf{a}_2 &= \frac{1}{\sqrt{3}}a \hat{\mathbf{y}} + \frac{1}{3}c \hat{\mathbf{z}} \\ \mathbf{a}_3 &= -\frac{1}{2}a \hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a \hat{\mathbf{y}} + \frac{1}{3}c \hat{\mathbf{z}}\end{aligned}$$





# Basis vectors

|                   | Lattice<br>coordinates                                                                                                                       |     | Cartesian<br>coordinates                                                                                                                                     | Wyckoff<br>position | Atom<br>type |
|-------------------|----------------------------------------------------------------------------------------------------------------------------------------------|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------|--------------|
| $\mathbf{B}_1$    | $= x_1 \mathbf{a}_1 + x_1 \mathbf{a}_2 + x_1 \mathbf{a}_3$                                                                                   | $=$ | $cx_1 \hat{\mathbf{z}}$                                                                                                                                      | (4c)                | Tl I         |
| $\mathbf{B}_2$    | $= -\left(x_1 - \frac{1}{2}\right) \mathbf{a}_1 - \left(x_1 - \frac{1}{2}\right) \mathbf{a}_2 - \left(x_1 - \frac{1}{2}\right) \mathbf{a}_3$ | $=$ | $-c\left(x_1 - \frac{1}{2}\right) \hat{\mathbf{z}}$                                                                                                          | (4c)                | Tl I         |
| $\mathbf{B}_3$    | $= -x_1 \mathbf{a}_1 - x_1 \mathbf{a}_2 - x_1 \mathbf{a}_3$                                                                                  | $=$ | $-cx_1 \hat{\mathbf{z}}$                                                                                                                                     | (4c)                | Tl I         |
| $\mathbf{B}_4$    | $= \left(x_1 + \frac{1}{2}\right) \mathbf{a}_1 + \left(x_1 + \frac{1}{2}\right) \mathbf{a}_2 + \left(x_1 + \frac{1}{2}\right) \mathbf{a}_3$  | $=$ | $c\left(x_1 + \frac{1}{2}\right) \hat{\mathbf{z}}$                                                                                                           | (4c)                | Tl I         |
| $\mathbf{B}_5$    | $= x_2 \mathbf{a}_1 - \left(x_2 - \frac{1}{2}\right) \mathbf{a}_2 + \frac{1}{4} \mathbf{a}_3$                                                | $=$ | $\frac{1}{8}a(4x_2 - 1) \hat{\mathbf{x}} - \frac{\sqrt{3}}{8}a(4x_2 - 1) \hat{\mathbf{y}} + \frac{1}{4}c \hat{\mathbf{z}}$                                   | (6e)                | Cl I         |
| $\mathbf{B}_6$    | $= \frac{1}{4} \mathbf{a}_1 + x_2 \mathbf{a}_2 - \left(x_2 - \frac{1}{2}\right) \mathbf{a}_3$                                                | $=$ | $\frac{1}{8}a(4x_2 - 1) \hat{\mathbf{x}} + \frac{\sqrt{3}}{8}a(4x_2 - 1) \hat{\mathbf{y}} + \frac{1}{4}c \hat{\mathbf{z}}$                                   | (6e)                | Cl I         |
| $\mathbf{B}_7$    | $= -\left(x_2 - \frac{1}{2}\right) \mathbf{a}_1 + \frac{1}{4} \mathbf{a}_2 + x_2 \mathbf{a}_3$                                               | $=$ | $-a\left(x_2 - \frac{1}{4}\right) \hat{\mathbf{x}} + \frac{1}{4}c \hat{\mathbf{z}}$                                                                          | (6e)                | Cl I         |
| $\mathbf{B}_8$    | $= -x_2 \mathbf{a}_1 + \left(x_2 + \frac{1}{2}\right) \mathbf{a}_2 + \frac{3}{4} \mathbf{a}_3$                                               | $=$ | $-\frac{1}{8}a(4x_2 + 3) \hat{\mathbf{x}} + \frac{\sqrt{3}}{24}a(12x_2 + 1) \hat{\mathbf{y}} + \frac{5}{12}c \hat{\mathbf{z}}$                               | (6e)                | Cl I         |
| $\mathbf{B}_9$    | $= \frac{3}{4} \mathbf{a}_1 - x_2 \mathbf{a}_2 + \left(x_2 + \frac{1}{2}\right) \mathbf{a}_3$                                                | $=$ | $-\frac{1}{8}a(4x_2 - 1) \hat{\mathbf{x}} - \frac{\sqrt{3}}{24}a(12x_2 + 5) \hat{\mathbf{y}} + \frac{5}{12}c \hat{\mathbf{z}}$                               | (6e)                | Cl I         |
| $\mathbf{B}_{10}$ | $= \left(x_2 + \frac{1}{2}\right) \mathbf{a}_1 + \frac{3}{4} \mathbf{a}_2 - x_2 \mathbf{a}_3$                                                | $=$ | $a\left(x_2 + \frac{1}{4}\right) \hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a \hat{\mathbf{y}} + \frac{5}{12}c \hat{\mathbf{z}}$                                   | (6e)                | Cl I         |
| $\mathbf{B}_{11}$ | $= x_3 \mathbf{a}_1 - \left(x_3 - \frac{1}{2}\right) \mathbf{a}_2 + \frac{1}{4} \mathbf{a}_3$                                                | $=$ | $\frac{1}{8}a(4x_3 - 1) \hat{\mathbf{x}} - \frac{\sqrt{3}}{8}a(4x_3 - 1) \hat{\mathbf{y}} + \frac{1}{4}c \hat{\mathbf{z}}$                                   | (6e)                | Cs I         |
| $\mathbf{B}_{12}$ | $= \frac{1}{4} \mathbf{a}_1 + x_3 \mathbf{a}_2 - \left(x_3 - \frac{1}{2}\right) \mathbf{a}_3$                                                | $=$ | $\frac{1}{8}a(4x_3 - 1) \hat{\mathbf{x}} + \frac{\sqrt{3}}{8}a(4x_3 - 1) \hat{\mathbf{y}} + \frac{1}{4}c \hat{\mathbf{z}}$                                   | (6e)                | Cs I         |
| $\mathbf{B}_{13}$ | $= -\left(x_3 - \frac{1}{2}\right) \mathbf{a}_1 + \frac{1}{4} \mathbf{a}_2 + x_3 \mathbf{a}_3$                                               | $=$ | $-a\left(x_3 - \frac{1}{4}\right) \hat{\mathbf{x}} + \frac{1}{4}c \hat{\mathbf{z}}$                                                                          | (6e)                | Cs I         |
| $\mathbf{B}_{14}$ | $= -x_3 \mathbf{a}_1 + \left(x_3 + \frac{1}{2}\right) \mathbf{a}_2 + \frac{3}{4} \mathbf{a}_3$                                               | $=$ | $-\frac{1}{8}a(4x_3 + 3) \hat{\mathbf{x}} + \frac{\sqrt{3}}{24}a(12x_3 + 1) \hat{\mathbf{y}} + \frac{5}{12}c \hat{\mathbf{z}}$                               | (6e)                | Cs I         |
| $\mathbf{B}_{15}$ | $= \frac{3}{4} \mathbf{a}_1 - x_3 \mathbf{a}_2 + \left(x_3 + \frac{1}{2}\right) \mathbf{a}_3$                                                | $=$ | $-\frac{1}{8}a(4x_3 - 1) \hat{\mathbf{x}} - \frac{\sqrt{3}}{24}a(12x_3 + 5) \hat{\mathbf{y}} + \frac{5}{12}c \hat{\mathbf{z}}$                               | (6e)                | Cs I         |
| $\mathbf{B}_{16}$ | $= \left(x_3 + \frac{1}{2}\right) \mathbf{a}_1 + \frac{3}{4} \mathbf{a}_2 - x_3 \mathbf{a}_3$                                                | $=$ | $a\left(x_3 + \frac{1}{4}\right) \hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a \hat{\mathbf{y}} + \frac{5}{12}c \hat{\mathbf{z}}$                                   | (6e)                | Cs I         |
| $\mathbf{B}_{17}$ | $= x_4 \mathbf{a}_1 + y_4 \mathbf{a}_2 + z_4 \mathbf{a}_3$                                                                                   | $=$ | $\frac{1}{2}a(x_4 - z_4) \hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a(x_4 - 2y_4 + z_4) \hat{\mathbf{y}} + \frac{1}{3}c(x_4 + y_4 + z_4) \hat{\mathbf{z}}$         | (12f)               | Cl II        |
| $\mathbf{B}_{18}$ | $= z_4 \mathbf{a}_1 + x_4 \mathbf{a}_2 + y_4 \mathbf{a}_3$                                                                                   | $=$ | $-\frac{1}{2}a(y_4 - z_4) \hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a(2x_4 - y_4 - z_4) \hat{\mathbf{y}} + \frac{1}{3}c(x_4 + y_4 + z_4) \hat{\mathbf{z}}$        | (12f)               | Cl II        |
| $\mathbf{B}_{19}$ | $= y_4 \mathbf{a}_1 + z_4 \mathbf{a}_2 + x_4 \mathbf{a}_3$                                                                                   | $=$ | $-\frac{1}{2}a(x_4 - y_4) \hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a(x_4 + y_4 - 2z_4) \hat{\mathbf{y}} + \frac{1}{3}c(x_4 + y_4 + z_4) \hat{\mathbf{z}}$        | (12f)               | Cl II        |
| $\mathbf{B}_{20}$ | $= -\left(z_4 - \frac{1}{2}\right) \mathbf{a}_1 - \left(y_4 - \frac{1}{2}\right) \mathbf{a}_2 - \left(x_4 - \frac{1}{2}\right) \mathbf{a}_3$ | $=$ | $\frac{1}{2}a(x_4 - z_4) \hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a(x_4 - 2y_4 + z_4) \hat{\mathbf{y}} - \frac{1}{6}c(2x_4 + 2y_4 + 2z_4 - 3) \hat{\mathbf{z}}$  | (12f)               | Cl II        |
| $\mathbf{B}_{21}$ | $= -\left(y_4 - \frac{1}{2}\right) \mathbf{a}_1 - \left(x_4 - \frac{1}{2}\right) \mathbf{a}_2 - \left(z_4 - \frac{1}{2}\right) \mathbf{a}_3$ | $=$ | $-\frac{1}{2}a(y_4 - z_4) \hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a(2x_4 - y_4 - z_4) \hat{\mathbf{y}} - \frac{1}{6}c(2x_4 + 2y_4 + 2z_4 - 3) \hat{\mathbf{z}}$ | (12f)               | Cl II        |
| $\mathbf{B}_{22}$ | $= -\left(x_4 - \frac{1}{2}\right) \mathbf{a}_1 - \left(z_4 - \frac{1}{2}\right) \mathbf{a}_2 - \left(y_4 - \frac{1}{2}\right) \mathbf{a}_3$ | $=$ | $-\frac{1}{2}a(x_4 - y_4) \hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a(x_4 + y_4 - 2z_4) \hat{\mathbf{y}} - \frac{1}{6}c(2x_4 + 2y_4 + 2z_4 - 3) \hat{\mathbf{z}}$ | (12f)               | Cl II        |
| $\mathbf{B}_{23}$ | $= -x_4 \mathbf{a}_1 - y_4 \mathbf{a}_2 - z_4 \mathbf{a}_3$                                                                                  | $=$ | $-\frac{1}{2}a(x_4 - z_4) \hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a(x_4 - 2y_4 + z_4) \hat{\mathbf{y}} - \frac{1}{3}c(x_4 + y_4 + z_4) \hat{\mathbf{z}}$        | (12f)               | Cl II        |
| $\mathbf{B}_{24}$ | $= -z_4 \mathbf{a}_1 - x_4 \mathbf{a}_2 - y_4 \mathbf{a}_3$                                                                                  | $=$ | $\frac{1}{2}a(y_4 - z_4) \hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a(2x_4 - y_4 - z_4) \hat{\mathbf{y}} - \frac{1}{3}c(x_4 + y_4 + z_4) \hat{\mathbf{z}}$         | (12f)               | Cl II        |
| $\mathbf{B}_{25}$ | $= -y_4 \mathbf{a}_1 - z_4 \mathbf{a}_2 - x_4 \mathbf{a}_3$                                                                                  | $=$ | $\frac{1}{2}a(x_4 - y_4) \hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a(x_4 + y_4 - 2z_4) \hat{\mathbf{y}} - \frac{1}{3}c(x_4 + y_4 + z_4) \hat{\mathbf{z}}$         | (12f)               | Cl II        |
| $\mathbf{B}_{26}$ | $= \left(z_4 + \frac{1}{2}\right) \mathbf{a}_1 + \left(y_4 + \frac{1}{2}\right) \mathbf{a}_2 + \left(x_4 + \frac{1}{2}\right) \mathbf{a}_3$  | $=$ | $-\frac{1}{2}a(x_4 - z_4) \hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a(x_4 - 2y_4 + z_4) \hat{\mathbf{y}} + \frac{1}{6}c(2x_4 + 2y_4 + 2z_4 + 3) \hat{\mathbf{z}}$ | (12f)               | Cl II        |
| $\mathbf{B}_{27}$ | $= \left(y_4 + \frac{1}{2}\right) \mathbf{a}_1 + \left(x_4 + \frac{1}{2}\right) \mathbf{a}_2 + \left(z_4 + \frac{1}{2}\right) \mathbf{a}_3$  | $=$ | $\frac{1}{2}a(y_4 - z_4) \hat{\mathbf{x}} + \frac{\sqrt{3}}{6}a(2x_4 - y_4 - z_4) \hat{\mathbf{y}} + \frac{1}{6}c(2x_4 + 2y_4 + 2z_4 + 3) \hat{\mathbf{z}}$  | (12f)               | Cl II        |
| $\mathbf{B}_{28}$ | $= \left(x_4 + \frac{1}{2}\right) \mathbf{a}_1 + \left(z_4 + \frac{1}{2}\right) \mathbf{a}_2 + \left(y_4 + \frac{1}{2}\right) \mathbf{a}_3$  | $=$ | $\frac{1}{2}a(x_4 - y_4) \hat{\mathbf{x}} - \frac{\sqrt{3}}{6}a(x_4 + y_4 - 2z_4) \hat{\mathbf{y}} + \frac{1}{6}c(2x_4 + 2y_4 + 2z_4 + 3) \hat{\mathbf{z}}$  | (12f)               | Cl II        |

## References

- [1] J. L. Hoard and L. Goldstein, *The Crystal Structure of Cesium Enneachlordithalliate,  $Cs_3Tl_2Cl_9$* , J. Chem. Phys. **3**, 199–202 (1935), doi:10.1063/1.1749633.
- [2] R. W. G. Wyckoff, *The Analytical Expression of the Results of the Theory of Space-Groups*, vol. 318 (Carnegie Institution of Washington, Washington DC, 1922).

## Found in

- [1] R. T. Downs and M. Hall-Wallace, *The American Mineralogist Crystal Structure Database*, Am. Mineral. **88**, 247–250 (2003).

## First cited in

D. Hicks, M. J. Mehl, M. Esters, C. Oses, O. Levy, G. L. W. Hart, C. Toher, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 3*, Comput. Mater. Sci. **199**, 110450 (2021), doi: 10.1016/j.commatsci.2021.110450.