

AsKSe₂ (*P*1) Structure: ABC2_aP16_1_4a_4a_8a-001


This structure previously used the label(s) **ABC2_aP16_1_4a_4a_8a**. Calls to these addresses will be redirected here.

Cite this page as:

- H. Eckert, S. Divilov, M. J. Mehl, D. Hicks, A. C. Zettel, M. Esters, X. Campilongo, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 4*, Comput. Mater. Sci. **240**, 112988 (2024), doi: 10.1016/j.commatsci.2024.112988.
- D. Hicks, M. J. Mehl, M. Esters, C. Oses, O. Levy, G. L. W. Hart, C. Toher, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 3*, Comput. Mater. Sci. **199**, 110450 (2021), doi: 10.1016/j.commatsci.2021.110450.

<https://aflow.org/prototype-encyclopedia/GZNE>


https://aflow.org/prototype-encyclopedia/ABC2_aP16_1_4a_4a_8a-001



Prototype	AsKSe ₂
AFLOW prototype label	ABC2_aP16_1_4a_4a_8a-001
ICSD	65297
CCDC	1627427
Pearson symbol	aP16
Space group number	1
Space group symbol	<i>P</i> 1
AFLOW prototype command	<pre>aflow --proto=ABC2_aP16_1_4a_4a_8a-001 --params=a,b/a,c/a,α,β,γ,x₁,y₁,z₁,x₂,y₂,z₂,x₃,y₃,z₃,x₄,y₄,z₄,x₅,y₅,z₅,x₆,y₆, z₆,x₇,y₇,z₇,x₈,y₈,z₈,x₉,y₉,z₉,x₁₀,y₁₀,z₁₀,x₁₁,y₁₁,z₁₁,x₁₂,y₁₂,z₁₂,x₁₃,y₁₃,z₁₃,x₁₄,y₁₄,z₁₄, x₁₅,y₁₅,z₁₅,x₁₆,y₁₆,z₁₆</pre>

Triclinic primitive vectors

$$\begin{aligned}
\mathbf{a}_1 &= a \hat{\mathbf{x}} \\
\mathbf{a}_2 &= b \cos \gamma \hat{\mathbf{x}} + b \sin \gamma \hat{\mathbf{y}} \\
\mathbf{a}_3 &= c_x \hat{\mathbf{x}} + c_y \hat{\mathbf{y}} + c_z \hat{\mathbf{z}} \\
c_x &= c \cos \beta \\
c_y &= c(\cos \alpha - \cos \beta \cos \gamma) / \sin \gamma \\
c_z &= \sqrt{c^2 - c_x^2 - c_y^2}
\end{aligned}$$



Basis vectors

	Lattice coordinates		Cartesian coordinates	Wyckoff position	Atom type
$\mathbf{B}_1$	$x_1 \mathbf{a}_1 + y_1 \mathbf{a}_2 + z_1 \mathbf{a}_3$	=	$(ax_1 + by_1 \cos \gamma + c_x z_1) \hat{\mathbf{x}} + (by_1 \sin \gamma + c_y z_1) \hat{\mathbf{y}} + c_z z_1 \hat{\mathbf{z}}$	(1a)	As I
$\mathbf{B}_2$	$x_2 \mathbf{a}_1 + y_2 \mathbf{a}_2 + z_2 \mathbf{a}_3$	=	$(ax_2 + by_2 \cos \gamma + c_x z_2) \hat{\mathbf{x}} + (by_2 \sin \gamma + c_y z_2) \hat{\mathbf{y}} + c_z z_2 \hat{\mathbf{z}}$	(1a)	As II
$\mathbf{B}_3$	$x_3 \mathbf{a}_1 + y_3 \mathbf{a}_2 + z_3 \mathbf{a}_3$	=	$(ax_3 + by_3 \cos \gamma + c_x z_3) \hat{\mathbf{x}} + (by_3 \sin \gamma + c_y z_3) \hat{\mathbf{y}} + c_z z_3 \hat{\mathbf{z}}$	(1a)	As III
$\mathbf{B}_4$	$x_4 \mathbf{a}_1 + y_4 \mathbf{a}_2 + z_4 \mathbf{a}_3$	=	$(ax_4 + by_4 \cos \gamma + c_x z_4) \hat{\mathbf{x}} + (by_4 \sin \gamma + c_y z_4) \hat{\mathbf{y}} + c_z z_4 \hat{\mathbf{z}}$	(1a)	As IV
$\mathbf{B}_5$	$x_5 \mathbf{a}_1 + y_5 \mathbf{a}_2 + z_5 \mathbf{a}_3$	=	$(ax_5 + by_5 \cos \gamma + c_x z_5) \hat{\mathbf{x}} + (by_5 \sin \gamma + c_y z_5) \hat{\mathbf{y}} + c_z z_5 \hat{\mathbf{z}}$	(1a)	K I
$\mathbf{B}_6$	$x_6 \mathbf{a}_1 + y_6 \mathbf{a}_2 + z_6 \mathbf{a}_3$	=	$(ax_6 + by_6 \cos \gamma + c_x z_6) \hat{\mathbf{x}} + (by_6 \sin \gamma + c_y z_6) \hat{\mathbf{y}} + c_z z_6 \hat{\mathbf{z}}$	(1a)	K II
$\mathbf{B}_7$	$x_7 \mathbf{a}_1 + y_7 \mathbf{a}_2 + z_7 \mathbf{a}_3$	=	$(ax_7 + by_7 \cos \gamma + c_x z_7) \hat{\mathbf{x}} + (by_7 \sin \gamma + c_y z_7) \hat{\mathbf{y}} + c_z z_7 \hat{\mathbf{z}}$	(1a)	K III
$\mathbf{B}_8$	$x_8 \mathbf{a}_1 + y_8 \mathbf{a}_2 + z_8 \mathbf{a}_3$	=	$(ax_8 + by_8 \cos \gamma + c_x z_8) \hat{\mathbf{x}} + (by_8 \sin \gamma + c_y z_8) \hat{\mathbf{y}} + c_z z_8 \hat{\mathbf{z}}$	(1a)	K IV
$\mathbf{B}_9$	$x_9 \mathbf{a}_1 + y_9 \mathbf{a}_2 + z_9 \mathbf{a}_3$	=	$(ax_9 + by_9 \cos \gamma + c_x z_9) \hat{\mathbf{x}} + (by_9 \sin \gamma + c_y z_9) \hat{\mathbf{y}} + c_z z_9 \hat{\mathbf{z}}$	(1a)	Se I
$\mathbf{B}_{10}$	$x_{10} \mathbf{a}_1 + y_{10} \mathbf{a}_2 + z_{10} \mathbf{a}_3$	=	$(ax_{10} + by_{10} \cos \gamma + c_x z_{10}) \hat{\mathbf{x}} + (by_{10} \sin \gamma + c_y z_{10}) \hat{\mathbf{y}} + c_z z_{10} \hat{\mathbf{z}}$	(1a)	Se II
$\mathbf{B}_{11}$	$x_{11} \mathbf{a}_1 + y_{11} \mathbf{a}_2 + z_{11} \mathbf{a}_3$	=	$(ax_{11} + by_{11} \cos \gamma + c_x z_{11}) \hat{\mathbf{x}} + (by_{11} \sin \gamma + c_y z_{11}) \hat{\mathbf{y}} + c_z z_{11} \hat{\mathbf{z}}$	(1a)	Se III
$\mathbf{B}_{12}$	$x_{12} \mathbf{a}_1 + y_{12} \mathbf{a}_2 + z_{12} \mathbf{a}_3$	=	$(ax_{12} + by_{12} \cos \gamma + c_x z_{12}) \hat{\mathbf{x}} + (by_{12} \sin \gamma + c_y z_{12}) \hat{\mathbf{y}} + c_z z_{12} \hat{\mathbf{z}}$	(1a)	Se IV
$\mathbf{B}_{13}$	$x_{13} \mathbf{a}_1 + y_{13} \mathbf{a}_2 + z_{13} \mathbf{a}_3$	=	$(ax_{13} + by_{13} \cos \gamma + c_x z_{13}) \hat{\mathbf{x}} + (by_{13} \sin \gamma + c_y z_{13}) \hat{\mathbf{y}} + c_z z_{13} \hat{\mathbf{z}}$	(1a)	Se V
$\mathbf{B}_{14}$	$x_{14} \mathbf{a}_1 + y_{14} \mathbf{a}_2 + z_{14} \mathbf{a}_3$	=	$(ax_{14} + by_{14} \cos \gamma + c_x z_{14}) \hat{\mathbf{x}} + (by_{14} \sin \gamma + c_y z_{14}) \hat{\mathbf{y}} + c_z z_{14} \hat{\mathbf{z}}$	(1a)	Se VI
$\mathbf{B}_{15}$	$x_{15} \mathbf{a}_1 + y_{15} \mathbf{a}_2 + z_{15} \mathbf{a}_3$	=	$(ax_{15} + by_{15} \cos \gamma + c_x z_{15}) \hat{\mathbf{x}} + (by_{15} \sin \gamma + c_y z_{15}) \hat{\mathbf{y}} + c_z z_{15} \hat{\mathbf{z}}$	(1a)	Se VII
$\mathbf{B}_{16}$	$x_{16} \mathbf{a}_1 + y_{16} \mathbf{a}_2 + z_{16} \mathbf{a}_3$	=	$(ax_{16} + by_{16} \cos \gamma + c_x z_{16}) \hat{\mathbf{x}} + (by_{16} \sin \gamma + c_y z_{16}) \hat{\mathbf{y}} + c_z z_{16} \hat{\mathbf{z}}$	(1a)	Se VIII

References

- [1] W. S. Sheldrick and H.-J. Häusler, *Zur Kenntnis von Alkalimetaselenoarseniten Darstellung und Kristallstrukturen von $MAsSe_2$, $M = K, Rb, Cs$* , Z. Anorganische und Allgemeine Chemie **561**, 139–148 (1988), doi:10.1002/zaac.19885610115.

Found in

- [1] P. Villars and L. Calvert, *Pearson's Handbook of Crystallographic Data for Intermetallic Phases* (ASM International, Materials Park, OH, 1991), 2nd edn.

First cited in

- M. J. Mehl, D. Hicks, C. Toher, O. Levy, R. M. Hanson, G. L. W. Hart, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 1*, Comput. Mater. Sci. **136**, S1-828 (2017). doi: 10.1016/j.commatsci.2017.01.017