

Mg[NH] Structure: ABC_hP36_175_jk_jk_jk-001


This structure previously used the label(s) `ABC_hP36_175_jk_jk_jk`. Calls to these addresses will be redirected here.

Cite this page as:

- H. Eckert, S. Divilov, M. J. Mehl, D. Hicks, A. C. Zettel, M. Esters, X. Campilongo, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 4*, Comput. Mater. Sci. **240**, 112988 (2024), doi: 10.1016/j.commatsci.2024.112988.
- D. Hicks, M. J. Mehl, M. Esters, C. Oses, O. Levy, G. L. W. Hart, C. Toher, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 3*, Comput. Mater. Sci. **199**, 110450 (2021), doi: 10.1016/j.commatsci.2021.110450.

<https://aflow.org/prototype-encyclopedia/12DH>

https://aflow.org/prototype-encyclopedia/ABC_hP36_175_jk_jk_jk-001




Prototype	HMgN
AFLOW prototype label	ABC_hP36_175_jk_jk_jk-001
ICSD	261304

CCDC	793340
Pearson symbol	hP36
Space group number	175
Space group symbol	$P6/m$
AFLOW prototype command	aflow --proto=ABC_hP36_175_jk_jk_jk-001 --params= $a, c/a, x_1, y_1, x_2, y_2, x_3, y_3, x_4, y_4, x_5, y_5, x_6, y_6$

- The experimental data was taken using deuterium.

Hexagonal primitive vectors

$$\begin{aligned}
\mathbf{a}_1 &= \frac{1}{2}a \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}a \hat{\mathbf{y}} \\
\mathbf{a}_2 &= \frac{1}{2}a \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}a \hat{\mathbf{y}} \\
\mathbf{a}_3 &= c \hat{\mathbf{z}}
\end{aligned}$$



Basis vectors

	Lattice coordinates		Cartesian coordinates	Wyckoff position	Atom type
$\mathbf{B}_1$	$= x_1 \mathbf{a}_1 + y_1 \mathbf{a}_2$	$=$	$\frac{1}{2}a(x_1 + y_1) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}a(x_1 - y_1) \hat{\mathbf{y}}$	(6j)	H I
$\mathbf{B}_2$	$= -y_1 \mathbf{a}_1 + (x_1 - y_1) \mathbf{a}_2$	$=$	$\frac{1}{2}a(x_1 - 2y_1) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}ax_1 \hat{\mathbf{y}}$	(6j)	H I
$\mathbf{B}_3$	$= -(x_1 - y_1) \mathbf{a}_1 - x_1 \mathbf{a}_2$	$=$	$-\frac{1}{2}a(2x_1 - y_1) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}ay_1 \hat{\mathbf{y}}$	(6j)	H I
$\mathbf{B}_4$	$= -x_1 \mathbf{a}_1 - y_1 \mathbf{a}_2$	$=$	$-\frac{1}{2}a(x_1 + y_1) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}a(x_1 - y_1) \hat{\mathbf{y}}$	(6j)	H I
$\mathbf{B}_5$	$= y_1 \mathbf{a}_1 - (x_1 - y_1) \mathbf{a}_2$	$=$	$\frac{1}{2}a(-x_1 + 2y_1) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}ax_1 \hat{\mathbf{y}}$	(6j)	H I
$\mathbf{B}_6$	$= (x_1 - y_1) \mathbf{a}_1 + x_1 \mathbf{a}_2$	$=$	$\frac{1}{2}a(2x_1 - y_1) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}ay_1 \hat{\mathbf{y}}$	(6j)	H I
$\mathbf{B}_7$	$= x_2 \mathbf{a}_1 + y_2 \mathbf{a}_2$	$=$	$\frac{1}{2}a(x_2 + y_2) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}a(x_2 - y_2) \hat{\mathbf{y}}$	(6j)	Mg I
$\mathbf{B}_8$	$= -y_2 \mathbf{a}_1 + (x_2 - y_2) \mathbf{a}_2$	$=$	$\frac{1}{2}a(x_2 - 2y_2) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}ax_2 \hat{\mathbf{y}}$	(6j)	Mg I
$\mathbf{B}_9$	$= -(x_2 - y_2) \mathbf{a}_1 - x_2 \mathbf{a}_2$	$=$	$-\frac{1}{2}a(2x_2 - y_2) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}ay_2 \hat{\mathbf{y}}$	(6j)	Mg I
$\mathbf{B}_{10}$	$= -x_2 \mathbf{a}_1 - y_2 \mathbf{a}_2$	$=$	$-\frac{1}{2}a(x_2 + y_2) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}a(x_2 - y_2) \hat{\mathbf{y}}$	(6j)	Mg I
$\mathbf{B}_{11}$	$= y_2 \mathbf{a}_1 - (x_2 - y_2) \mathbf{a}_2$	$=$	$\frac{1}{2}a(-x_2 + 2y_2) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}ax_2 \hat{\mathbf{y}}$	(6j)	Mg I
$\mathbf{B}_{12}$	$= (x_2 - y_2) \mathbf{a}_1 + x_2 \mathbf{a}_2$	$=$	$\frac{1}{2}a(2x_2 - y_2) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}ay_2 \hat{\mathbf{y}}$	(6j)	Mg I
$\mathbf{B}_{13}$	$= x_3 \mathbf{a}_1 + y_3 \mathbf{a}_2$	$=$	$\frac{1}{2}a(x_3 + y_3) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}a(x_3 - y_3) \hat{\mathbf{y}}$	(6j)	N I
$\mathbf{B}_{14}$	$= -y_3 \mathbf{a}_1 + (x_3 - y_3) \mathbf{a}_2$	$=$	$\frac{1}{2}a(x_3 - 2y_3) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}ax_3 \hat{\mathbf{y}}$	(6j)	N I
$\mathbf{B}_{15}$	$= -(x_3 - y_3) \mathbf{a}_1 - x_3 \mathbf{a}_2$	$=$	$-\frac{1}{2}a(2x_3 - y_3) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}ay_3 \hat{\mathbf{y}}$	(6j)	N I
$\mathbf{B}_{16}$	$= -x_3 \mathbf{a}_1 - y_3 \mathbf{a}_2$	$=$	$-\frac{1}{2}a(x_3 + y_3) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}a(x_3 - y_3) \hat{\mathbf{y}}$	(6j)	N I
$\mathbf{B}_{17}$	$= y_3 \mathbf{a}_1 - (x_3 - y_3) \mathbf{a}_2$	$=$	$\frac{1}{2}a(-x_3 + 2y_3) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}ax_3 \hat{\mathbf{y}}$	(6j)	N I
$\mathbf{B}_{18}$	$= (x_3 - y_3) \mathbf{a}_1 + x_3 \mathbf{a}_2$	$=$	$\frac{1}{2}a(2x_3 - y_3) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}ay_3 \hat{\mathbf{y}}$	(6j)	N I
$\mathbf{B}_{19}$	$= x_4 \mathbf{a}_1 + y_4 \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$\frac{1}{2}a(x_4 + y_4) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}a(x_4 - y_4) \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	H II
$\mathbf{B}_{20}$	$= -y_4 \mathbf{a}_1 + (x_4 - y_4) \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$\frac{1}{2}a(x_4 - 2y_4) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}ax_4 \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	H II
$\mathbf{B}_{21}$	$= -(x_4 - y_4) \mathbf{a}_1 - x_4 \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$-\frac{1}{2}a(2x_4 - y_4) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}ay_4 \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	H II

$\mathbf{B}_{22} =$	$-x_4 \mathbf{a}_1 - y_4 \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$-\frac{1}{2}a(x_4 + y_4) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}a(x_4 - y_4) \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	H II
$\mathbf{B}_{23} =$	$y_4 \mathbf{a}_1 - (x_4 - y_4) \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$\frac{1}{2}a(-x_4 + 2y_4) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}ax_4 \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	H II
$\mathbf{B}_{24} =$	$(x_4 - y_4) \mathbf{a}_1 + x_4 \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$\frac{1}{2}a(2x_4 - y_4) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}ay_4 \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	H II
$\mathbf{B}_{25} =$	$x_5 \mathbf{a}_1 + y_5 \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$\frac{1}{2}a(x_5 + y_5) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}a(x_5 - y_5) \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	Mg II
$\mathbf{B}_{26} =$	$-y_5 \mathbf{a}_1 + (x_5 - y_5) \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$\frac{1}{2}a(x_5 - 2y_5) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}ax_5 \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	Mg II
$\mathbf{B}_{27} =$	$-(x_5 - y_5) \mathbf{a}_1 - x_5 \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$-\frac{1}{2}a(2x_5 - y_5) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}ay_5 \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	Mg II
$\mathbf{B}_{28} =$	$-x_5 \mathbf{a}_1 - y_5 \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$-\frac{1}{2}a(x_5 + y_5) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}a(x_5 - y_5) \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	Mg II
$\mathbf{B}_{29} =$	$y_5 \mathbf{a}_1 - (x_5 - y_5) \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$\frac{1}{2}a(-x_5 + 2y_5) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}ax_5 \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	Mg II
$\mathbf{B}_{30} =$	$(x_5 - y_5) \mathbf{a}_1 + x_5 \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$\frac{1}{2}a(2x_5 - y_5) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}ay_5 \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	Mg II
$\mathbf{B}_{31} =$	$x_6 \mathbf{a}_1 + y_6 \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$\frac{1}{2}a(x_6 + y_6) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}a(x_6 - y_6) \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	N II
$\mathbf{B}_{32} =$	$-y_6 \mathbf{a}_1 + (x_6 - y_6) \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$\frac{1}{2}a(x_6 - 2y_6) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}ax_6 \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	N II
$\mathbf{B}_{33} =$	$-(x_6 - y_6) \mathbf{a}_1 - x_6 \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$-\frac{1}{2}a(2x_6 - y_6) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}ay_6 \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	N II
$\mathbf{B}_{34} =$	$-x_6 \mathbf{a}_1 - y_6 \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$-\frac{1}{2}a(x_6 + y_6) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}a(x_6 - y_6) \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	N II
$\mathbf{B}_{35} =$	$y_6 \mathbf{a}_1 - (x_6 - y_6) \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$\frac{1}{2}a(-x_6 + 2y_6) \hat{\mathbf{x}} - \frac{\sqrt{3}}{2}ax_6 \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	N II
$\mathbf{B}_{36} =$	$(x_6 - y_6) \mathbf{a}_1 + x_6 \mathbf{a}_2 + \frac{1}{2} \mathbf{a}_3$	$=$	$\frac{1}{2}a(2x_6 - y_6) \hat{\mathbf{x}} + \frac{\sqrt{3}}{2}ay_6 \hat{\mathbf{y}} + \frac{1}{2}c \hat{\mathbf{z}}$	(6k)	N II

References

- [1] F. Dolci, E. Napolitano, E. Weidner, S. Enzo, P. Moretto, M. Brunelli, T. Hansen, M. Fichtner, and W. Lohstroh, *Magnesium Imide: Synthesis and Structure Determination of an Unconventional Alkaline Earth Imide from Decomposition of Magnesium Amide*, Inorg. Chem. **50**, 1116–1122 (2001), doi:10.1021/ic1023778.

Found in

- [1] P. Villars and K. Cenzual, *Pearson's Crystal Data – Crystal Structure Database for Inorganic Compounds* (2013). ASM International.

First cited in

D. Hicks, M. J. Mehl, E. Gossett, C. Toher, O. Levy, R. M. Hanson, G. Hart, and S. Curtarolo, *The AFLOW Library of Crystallographic Prototypes: Part 2*, Comput. Mater. Sci. **161**, S1 (2019). doi: 10.1016/j.commatsci.2018.10.043